

GROTHENDIECK RINGS OF LAURENT SERIES FIELDS

RAF CLUCKERS*

ABSTRACT. We study Grothendieck rings (in the sense of logic) of fields. We prove the triviality of the Grothendieck rings of certain fields by constructing definable bijections which imply the triviality. More precisely, we consider valued fields, for example, fields of Laurent series over the real numbers, over p -adic numbers and over finite fields, and construct definable bijections from the line to the line minus one point.

1. INTRODUCTION

Recently, the Grothendieck ring of a structure, in the sense of logic, has been introduced in [3] and independently in [4]. The Grothendieck ring of a model-theoretical structure is built up as a quotient of the definable sets by definable bijections (see below), and thus, depends both on the model and the language. For $(M, \mathcal{L})$ a structure with the signature of a language $\mathcal{L}$ we write $K_0(M, \mathcal{L})$ for the Grothendieck ring of $(M, \mathcal{L})$. In [1] and [7], the following explicit calculations of Grothendieck rings of fields are made:

$K_0(\mathbb{R}, \mathcal{L}_{\text{ring}})$ is isomorphic to $\mathbb{Z}$,
 $K_0(\mathbb{Q}_p, \mathcal{L}_{\text{ring}})$ is trivial,
 $K_0(\mathbb{F}_p((t)), \mathcal{L}_{\text{ring}})$ is trivial.

Here, $\mathcal{L}_{\text{ring}}$ is the language $(+, -, \cdot, 0, 1)$. In [3] and [4] it is shown that the Grothendieck ring $K_0(\mathbb{C}, \mathcal{L}_{\text{ring}})$ is extremely big and complicated; $K_0(\mathbb{C}, \mathcal{L}_{\text{ring}})$, and many other Grothendieck rings, are not explicitly known.

Any Euler characteristic (in the sense of [1] or [4]), defined on the definable sets, factors through the natural projection of definable sets into the Grothendieck ring, and, in this sense, to know a Grothendieck ring is to know a universal Euler characteristic. Nevertheless, it happens that a Grothendieck ring is trivial.

The triviality of a Grothendieck ring can be proven by constructing a definable bijection from X to $X \setminus \{a\}$, where X is a definable set and $\{a\}$ a point on X . We develop general techniques to obtain definable bijections $K \rightarrow K^\times$, where K is a valued field and $K^\times = K \setminus \{0\}$. In section 1.2 we explain iterated Laurent series fields. In the present paper we prove:

Theorem 1. *Let L be either $\mathbb{Q}_p$ or a finite field extension of $\mathbb{Q}_p$, and let K be one of the fields L , $L((t_1))$, $L((t_1))((t_2))$, $L((t_1))((t_2))((t_3))$, and so on. Then $K_0(K, \mathcal{L}_{\text{ring}}) = 0$ and there exists a $\mathcal{L}_{\text{ring}}$ -definable bijection $K \rightarrow K^\times$.¹²*

2000 *Mathematics Subject Classification.* Primary 03C60, 12L12; Secondary 03C07.

Key words and phrases. Grothendieck rings, model theory, valued fields, Henselian rings.

*Research Assistant of the Fund for Scientific Research – Flanders (Belgium)(F.W.O.).

¹Here, as always, definable means definable with parameters.

²For $K = \mathbb{Q}_p$, this result was proven in [1].

Theorem 2. *Let L be $\mathbb{F}_q$ where $\mathbb{F}_q$ is the finite field with $q = p^l$ elements, p a prime, and let K be one of the fields $L((t_1))$, $L((t_1))((t_2))$, $L((t_1))((t_2))((t_3))$, and so on. Then $K_0(K, \mathcal{L}_{\text{ring}}) = 0$ and there exists a $\mathcal{L}_{\text{ring}}$ -definable bijection $K \rightarrow K^\times$.³*

Central in the proofs of this paper is a subgroup $H(K, \mathcal{L})$ of $\mathbb{Z}$, associated to a field K and a language $\mathcal{L}$, which is sensitive to some elementary arithmetical properties of the indices of n -th powers in $K^\times$ and of the number of n -th roots in $K^\times$ (see section 3). Using the definition, it follows immediately that, for example,

$$\begin{aligned} H(\mathbb{R}, \mathcal{L}_{\text{ring}}) &\text{ is } \mathbb{Z}, \\ H(\mathbb{Q}_p, \mathcal{L}_{\text{ring}}) &\text{ is } \mathbb{Z}, \text{ and} \\ H(\mathbb{C}, \mathcal{L}_{\text{ring}}) &\text{ is } \{0\}. \end{aligned}$$

We give two criteria for valued fields, for which the value group has a well-determined minimal strictly positive element, to have a trivial Grothendieck ring (proposition 1 and 2).

We also consider Laurent series fields over $\mathbb{R}$ and over fields of characteristic $p > 0$, using the language of Denef - Pas. The language of Denef - Pas [5] was introduced to study uniform p -adic integrals for all primes p , and is now still used in, for example, the theory of motivic integration (see [3] and [2]).

For any $\mathbb{Z}$ -valued field K with angular component map ac , the Grothendieck ring $K_0(K, \mathcal{L}_{\text{Pas}})$ is trivial, and there exists a $\mathcal{L}_{\text{Pas}}$ -definable bijection from K^2 onto $K^2 \setminus \{(0, 0)\}$, see [1], Thm. 1 and proposition 2 below. (Proposition 2 is more general than [1], Thm. 1.) The following theorems give stronger results for iterated Laurent series fields.

Theorem 3. *Let K be one of the fields $\mathbb{R}((t_1))$, $\mathbb{R}((t_1))((t_2))$, $\mathbb{R}((t_1))((t_2))((t_3))$, and so on. We have*

$$H(K, \mathcal{L}_{\text{ring}}) = \mathbb{Z}.$$

Endow K with a valuation onto a group of the form $\mathbb{Z}^k$ with lexicographical order, $k > 0$, and with the natural angular component map (as in section 2). Then

$$K_0(K, \mathcal{L}_{\text{Pas}}) = \{0\},$$

and there exist a bijection $K \rightarrow K^\times$, definable in the language $\mathcal{L}_{\text{Pas}}$ of Denef - Pas.

Theorem 4. *Let L be an arbitrary field of characteristic $p > 0$. Let K be one of the fields $L((t_1))$, $L((t_1))((t_2))$, $L((t_1))((t_2))((t_3))$ and so on. Endow K with a valuation onto a group of the form $\mathbb{Z}^k$ with lexicographical order, $k > 0$, and with the natural angular component map (as in section 2). Then $K_0(K, \mathcal{L}_{\text{Pas}}) = \{0\}$ and there exists a bijection $K \rightarrow K^\times$ definable in the language of Denef - Pas.*

1.1. Valued fields. Fix a field K . We call K a valued field if there is an ordered group $(G, +, \leq)$ and a valuation map $v : K \rightarrow G \cup \{\infty\}$ such that

- (i) $v(x) = \infty$ if and only if $x = 0$;
- (ii) $v(xy) = v(x) + v(y)$ for all $x, y \in K$;
- (iii) $v(x + y) \geq \min\{v(x), v(y)\}$ for all $x, y \in K$.

We write R for the valuation ring $\{x \in K \mid v(x) \geq 0\}$ of K , M for its unique maximal ideal and we write k for the residue field R/M and $p : R \rightarrow k$ for the natural projection. If $G = \mathbb{Z}$ we call K a $\mathbb{Z}$ -valued field. Whenever K is a $\mathbb{Z}$ -valued

³For $K = \mathbb{F}_q((t))$ this was proven in [1].

field, the valuation ring R is a discrete valuation ring, and a generator π of the maximal ideal of R is called a uniformizer.

A valued field often carries an angular component map modulo M , or angular component map for short; it is a group homomorphism $\text{ac} : K^\times \rightarrow k^\times$, extended by putting $\text{ac}(0) = 0$, and satisfying $\text{ac}(x) = p(x)$ for all x with $v(x) = 0$ (see [6]).

1.2. Iterated Laurent series fields. We define iterated Laurent series fields by induction. Let $L((t_1))$ be the field of (formal) Laurent series in the variable t_1 over L and let $L((t_1)) \dots ((t_{n-1}))((t_n))$ be the field of (formal) Laurent series in the variable t_n over $L((t_1)) \dots ((t_{n-1}))$. On a field $L((t_1)) \dots ((t_n))$ we can put many valuations, for example the valuation v_n taking values in the lexicographically ordered n -fold product of $\mathbb{Z}$, defined as follows. If $n = 1$, then we put $v_1(x) = s \in \mathbb{Z}$ whenever $x = \sum_{i \geq s} a_i t_1^i$ with $a_s \neq 0$ and $a_i \in L$. For general n , and $x = \sum_{i \geq s} a_i t_n^i$, where $a_s \neq 0$ and $a_i \in L((t_1)) \dots ((t_{n-1}))$, we put $v_n(x) = (s, v_{n-1}(a_s)) \in \mathbb{Z}^n$. Remark that the valuation ring with respect to the valuation v_n is Henselian.

1.3. Grothendieck rings. Let $\mathcal{L}$ be a language and let M be a model for $\mathcal{L}$. For $\mathcal{L}$ -definable sets $X \subset M^m$, $Y \subset M^n$, $m, n > 0$, a $\mathcal{L}$ -definable bijection $X \rightarrow Y$ is called an $\mathcal{L}$ -isomorphism and we write $X \cong_{\mathcal{L}} Y$, or $X \cong Y$ if the context is clear, if X and Y are $\mathcal{L}$ -isomorphic. (Definable always means definable with parameters.) For definable X and Y , we can choose disjoint definable sets $X', Y' \subset K^{m'}$ for some $m' > 0$, such that $X \cong X'$ and $Y \cong Y'$, and then we define the *disjoint union* $X \sqcup Y$ of X and Y up to isomorphism as $X' \cup Y'$. By the Grothendieck group $K_0(M, \mathcal{L})$ of the structure $(M, \mathcal{L})$ we mean the group generated by symbols $[A]$, for A a $\mathcal{L}$ -definable set, with the relations $[A] = [A']$ if $A \cong_{\mathcal{L}} A'$ and $[A] = [B] + [C]$ if A is the disjoint union of B and C . The group $K_0(M, \mathcal{L})$ carries a multiplicative structure induced by $[A \times B] = [A][B]$, where $A \times B$ is the Cartesian product of definable sets. The so-obtained ring is called the Grothendieck ring and for a $\mathcal{L}$ -definable set X we write $[X]$ for the image of X in $K_0(M, \mathcal{L})$.

Let T be a theory in some language $\mathcal{L}$. A formula φ with free variables $x_1, \dots, x_n$ determines a set in M^n for any model M of T . On these sets we can define a disjoint union operation and Cartesian products in the natural way. The Grothendieck group $K_0(T, \mathcal{L})$ is the group generated by symbols $[\varphi]$, for φ a $\mathcal{L}$ -formula, with the relations $[\varphi] = [\varphi']$ if the theory T implies that there is some $\mathcal{L}$ -definable bijection between the sets defined by φ and φ' , and the relation $[\varphi] = [\psi] + [\psi']$ if φ is the disjoint union of ψ and ψ' . This group carries a multiplicative structure induced by the Cartesian product of definable sets and the so obtained ring is called the Grothendieck ring of $(T, \mathcal{L})$.

2. LANGUAGES OF DENEUF - PAS

Let K be a valued field, with a valuation map $v : K \rightarrow G \cup \{\infty\}$ for some ordered group G , and an angular component map $\text{ac} : K \rightarrow k$, where k is the residue field. Let $\mathcal{L}_k$ be an arbitrary expansion of $\mathcal{L}_{\text{ring}}$ and let $\mathcal{L}_G$ be an arbitrary expansion of the language of ordered groups with infinity, namely $(+, -, 0, \infty, \leq)$. A language of Denef - Pas can in fact be either language in a wide variety of languages; it is always a three-sorted language of the form $(\mathcal{L}_k, \mathcal{L}_{\text{ring}}, \mathcal{L}_G, v, \text{ac})$, with as sorts:

- (i) a k -sort for the residue field-sort,
- (ii) a K -sort for the valued field-sort, and
- (iii) a G -sort for the value group-sort.

The language $\mathcal{L}_{\text{ring}}$ is used for the K -sort, $\mathcal{L}_k$ for the k -sort and $\mathcal{L}_G$ for the G -sort. The function symbol v stands for the valuation map $K \rightarrow G \cup \{\infty\}$ and ac stands for an angular component map $K \rightarrow k$ (in fact, this is an angular component map modulo the maximal ideal M). A structure for a language of Denef - Pas is denoted $(k, K, G \cup \{\infty\})$, where k, K and G are as above.

Remark that if $G = \mathbb{Z}$, namely if K is a $\mathbb{Z}$ -valued field, there exists a natural angular component map $\text{ac} : K \rightarrow k$ sending $x \neq 0$ to $t^{-v(x)}x \bmod M$, where t is a uniformizer of the valuation ring. More generally, if the value group of K is $\mathbb{Z}^n$, and $t_1, \dots, t_n$ are field elements such that $v(t_1) = (1, 0, \dots, 0), \dots, v(t_n) = (0, \dots, 0, 1)$ forms a set of generators of $\mathbb{Z}^n$, there is a natural angular component map $\text{ac} : K \rightarrow k$ given by $\text{ac}(x) = x \prod_i t_i^{-r_i} \bmod M$, where $v(x) = (r_1, \dots, r_n)$. These angular component maps are canonical up to the choice of t and t_i . Languages of Denef - Pas are denoted $\mathcal{L}_{\text{Pas}}$.

3. CALCULATIONS OF GROTHENDIECK RINGS

Let K be a field and $\mathcal{L}$ an expansion of $\mathcal{L}_{\text{ring}}$. We write $P_n(K)$ or P_n for the n -th powers in $K^\times$. For $n > 1$ we put

$$r_n(K) = \#\{x \in K \mid x^n = 1\}$$

and

$$s_n(K) = [K^\times : P_n(K)]$$

which is either a nonnegative integer or ∞ .

Definition 1. Let K be a field and let the numbers $s_n(K)$ and $r_n(K)$, $n > 1$ be as above. For $n > 1$ we put

$$\lambda_n(K, \mathcal{L}) = \frac{s_n(K)}{r_n(K)}$$

if the following conditions are satisfied

- $s_n(K) < \infty$ and $\frac{s_n(K)}{r_n(K)} \in \mathbb{Z}$;
- there exists a $\mathcal{L}$ -definable n -th root function. This means that there exists a definable set $\sqrt[n]{P_n}$ and a definable bijection $\sqrt[n]{\cdot} : P_n(K) \rightarrow \sqrt[n]{P_n}$ such that $(\sqrt[n]{x})^n = x$ for each $x \in P_n(K)$.

If one of the above conditions is not satisfied, we put $\lambda_n(K, \mathcal{L}) = 1$. We define $H(K, \mathcal{L})$ as the subgroup of $\mathbb{Z}$ generated by the numbers

$$\lambda_n(K, \mathcal{L}) - 1$$

for all $n > 1$.

Remark that if $\mathcal{L}'$ is an expansion of $\mathcal{L}$, then there is a group inclusion $H(K, \mathcal{L}) \rightarrow H(K, \mathcal{L}')$. Let $\mathcal{L}_v = (\mathcal{L}_{\text{ring}}, R)$ be the language of rings with an extra 1-ary relation symbol R which corresponds to a valuation ring inside the model. If the model is a valued field, we take the natural interpretations.

Lemma 1. Let K be a field and $\mathcal{L}$ an expansion of $\mathcal{L}_{\text{ring}}$. For each positive number $m \in H(K, \mathcal{L})$, there exists a $\mathcal{L}$ -definable bijection

$$\bigsqcup_{i=1}^{m+1} K^\times \rightarrow K^\times,$$

and thus, in $K_0(K, \mathcal{L})$,

$$m[K^\times] = 0.$$

Moreover, if K is a valued field and $\mathcal{L}$ is an expansion of $\mathcal{L}_v$, there exists a $\mathcal{L}$ -definable bijection

$$\bigsqcup_{i=1}^{m+1} (R \setminus \{0\}) \rightarrow R \setminus \{0\},$$

and thus, in $K_0(K, \mathcal{L})$,

$$m[R \setminus \{0\}] = 0.$$

Proof. We first prove that $K^\times \cong \bigsqcup_{i=1}^{\lambda_n} K^\times$ for all $\lambda_n = \lambda_n(K, \mathcal{L})$, $n > 1$. If $\lambda_n = 1$ there is nothing to prove, so let $\lambda_n > 1$. Remark that for each $x \in K^\times$ and each definable set $A \subset K$ there is a definable bijection $xA \cong A$. With the notation of definition 1, the sets $x\sqrt[n]{P_n}$ form a partition of $K^\times$ when x runs over the n -th roots of unity. This gives $\bigsqcup_{i=1}^{r_n} \sqrt[n]{P_n} \cong K^\times$. Since $K^\times$ is the disjoint union of all cosets of P_n inside $K^\times$, we find $\bigsqcup_{i=1}^{s_n} P_n \cong K^\times$. Combining with the isomorphism $P_n \cong \sqrt[n]{P_n}$ we calculate:

$$K^\times \cong \bigsqcup_{i=1}^{s_n} P_n \cong \bigsqcup_{i=1}^{s_n} \sqrt[n]{P_n} \cong \bigsqcup_{i=1}^{\lambda_n} \left(\bigsqcup_{i=1}^{r_n} \sqrt[n]{P_n} \right) \cong \bigsqcup_{i=1}^{\lambda_n} K^\times,$$

where $s_n = s_n(K)$ and $r_n = r_n(K)$. Now let $m > 0$ be in $H(K, \mathcal{L})$ and let $n > 1$, $s > 0$ be integers. By what we just have shown, we can add $\lambda_n - 1$ disjoint copies of $K^\times$ to $\bigsqcup_{i=1}^s K^\times$, in the sense that $\bigsqcup_{i=1}^s K^\times \cong \bigsqcup_{i=1}^{s+\lambda_n-1} K^\times$. Similarly, if $s > \lambda_n - 1$, we can subtract $\lambda_n - 1$ disjoint copies of $K^\times$ from $\bigsqcup_{i=1}^s K^\times$, to be precise, $\bigsqcup_{i=1}^s K^\times \cong \bigsqcup_{i=1}^{s-\lambda_n+1} K^\times$. The lemma follows since the numbers $\lambda_n - 1$ generate $H(K, \mathcal{L})$.

If $\mathcal{L}$ is an expansion of $\mathcal{L}_v$, we have the same isomorphisms and the same arguments for $R \setminus \{0\}$ instead of $K^\times$, working with $R \cap P_n$ and $R \cap \sqrt[n]{P_n}$ instead of P_n and $\sqrt[n]{P_n}$. $\square$

Definition 2. Let R be a valuation ring such that the value group has a minimal strictly positive element. Let $\pi \in R$ have minimal strictly positive valuation. Write M for the maximal ideal of R . Let ac be an angular component map $K \rightarrow k$, where k is the residue field. We define the set $R^{(1)}$ as

$$R^{(1)} = \{x \in R \mid \text{ac}(x) = 1\}.$$

The set $R^{(1)}$ is not necessarily definable in the language $\mathcal{L}_{\text{ring}}$. If $R^{(1)}$ is definable in some language $\mathcal{L}$ we have the following criterion. Remark also that a minimal strictly positive element in the value group necessarily is unique.

Proposition 1. Let K be a valued field. Suppose that the value group has a minimal strictly positive element and let $\pi \in R$ have this minimal strictly positive valuation. Let $\mathcal{L}$ be an expansion of $\mathcal{L}_v$ and let ac be an angular component map $K \rightarrow k$, with k the residue field. If $R^{(1)}$ is $\mathcal{L}$ -definable and $H(K, \mathcal{L}) = \mathbb{Z}$, then

$$K_0(K, \mathcal{L}) = 0, \quad K \cong_{\mathcal{L}} K^\times, \quad \text{and} \quad R \cong_{\mathcal{L}} R \setminus \{0\}.$$

Proof. We first prove that $K_0(K, \mathcal{L}) = 0$. We may suppose that $\text{ac}(\pi) = 1$, otherwise we could replace π by π/a where a is an arbitrary element with $v(a) = 0$ and $\text{ac}(a) = \text{ac}(\pi)$. The following is a $\mathcal{L}$ -isomorphism

$$R \sqcup R^{(1)} \rightarrow R^{(1)} : \begin{cases} x \in R & \mapsto 1 + \pi x, \\ x \in R^{(1)} & \mapsto \pi x. \end{cases}$$

This implies, in $K_0(K, \mathcal{L})$, that $[R] + [R^{(1)}] = [R^{(1)}]$, and thus after cancellation, $[R] = 0$. By lemma 1 and because $H(K, \mathcal{L}) = \mathbb{Z}$, also $[R \setminus \{0\}] = 0$. The following calculation implies $K_0(K, \mathcal{L}) = 0$:

$$0 = [R] = [R \setminus \{0\}] + [\{0\}] = [\{0\}] = 1.$$

We have $[\{0\}] = 1$ because $[\{0\}]$ is the multiplicative unit in $K_0(K, \mathcal{L})$.

Next we prove $R \cong R \setminus \{0\}$, by taking translates and applying homotheties to the occurring sets. We make all occurring disjoint unions explicit. Write f_1 for the isomorphism

$$f_1 : 1 + \pi^2(R \setminus \{0\}) \rightarrow \pi^2(R \setminus \{0\}) \cup 1 + \pi^2(R \setminus \{0\}),$$

given by lemma 1, it is an isomorphism from one copy of $R \setminus \{0\}$ onto two disjoint copies of $R \setminus \{0\}$. Define the function f_2 on $\pi^2 R \cup \pi + \pi^2 R^{(1)}$ by

$$f_2 : \pi^2 R \cup \pi + \pi^2 R^{(1)} \rightarrow \pi + \pi^2 R^{(1)} : \begin{cases} \pi^2 x & \mapsto \pi + \pi^2(1 + \pi x), \\ \pi + \pi^2 x & \mapsto \pi + \pi^2(\pi x), \end{cases}$$

then f_2 is an isomorphism from the disjoint union of R and $R^{(1)}$ to a copy of $R^{(1)}$. Finally, we find $\mathcal{L}$ -isomorphisms:

$$f : R \rightarrow R \setminus \{0\} : x \mapsto \begin{cases} f_1(x) & \text{if } x \in 1 + \pi^2(R \setminus \{0\}), \\ f_2(x) & \text{if } x \in \pi^2 R \cup \pi + \pi^2 R^{(1)}, \\ x & \text{else} \end{cases}$$

and

$$K \rightarrow K^\times : x \mapsto \begin{cases} f(x) & \text{if } x \in R, \\ x & \text{else.} \end{cases}$$

□

Proposition 1 immediately yields the triviality of the Grothendieck rings of $\mathbb{Q}_p$ and of $\mathbb{F}_q((t))$ with characteristic different from 2, which was originally proven in [1]. Theorems 1 and 2 of the present paper are generalizations. In case that $H(K, \mathcal{L})$ is different from $\mathbb{Z}$, we formulate the following criterion. (The argument of this criterion is similar to the proof of [1], Thm. 1.)

Proposition 2. *Let K be a valued field. Suppose that the value group has a unique minimal, strictly positive element and let $\pi \in R$ have this minimal strictly positive valuation. Let $\mathcal{L}$ be an expansion of $\mathcal{L}_v$ and let ac be an angular component map $K \rightarrow k$. If $R^{(1)}$ is $\mathcal{L}$ -definable, then*

$$K_0(K, \mathcal{L}) = 0, \quad K^2 \cong_{\mathcal{L}} K^2 \setminus \{(0, 0)\}, \quad \text{and} \quad R^2 \cong_{\mathcal{L}} R^2 \setminus \{(0, 0)\}.$$

Proof. We first prove that $K_0(K, \mathcal{L}) = 0$. As above we may suppose that $\text{ac}(\pi) = 1$. The following is a $\mathcal{L}$ -isomorphism

$$g_1 : R \sqcup R^{(1)} \rightarrow R^{(1)} : \begin{cases} x \in R & \mapsto 1 + \pi x, \\ x \in R^{(1)} & \mapsto \pi x. \end{cases}$$

As above, this implies that $[R] = 0$ in $K_0(K, \mathcal{L})$.

We argue that the disjoint union of two copies of $(R \setminus \{0\})^2$ is $\mathcal{L}$ -isomorphic to $(R \setminus \{0\})^2$ itself. Define the sets

$$\begin{aligned} X_1 &= \{(x, y) \in (R \setminus \{0\})^2 \mid v(x) \leq v(y)\}, \\ X_2 &= \{(x, y) \in (R \setminus \{0\})^2 \mid v(x) > v(y)\}, \end{aligned}$$

then X_1, X_2 form a partition of $(R \setminus \{0\})^2$. The isomorphisms

$$\begin{aligned} (R \setminus \{0\})^2 &\rightarrow X_1 : (x, y) \mapsto (x, xy), \\ (R \setminus \{0\})^2 &\rightarrow X_2 : (x, y) \mapsto (\pi xy, y), \end{aligned}$$

imply that $(R \setminus \{0\})^2 \sqcup (R \setminus \{0\})^2$ is isomorphic to $X_1 \cup X_2$ which is exactly $(R \setminus \{0\})^2$. After cancellation, it follows that $[(R \setminus \{0\})^2] = 0$.

Since $0 = [R] = [R \setminus \{0\}] + [\{0\}] = [R \setminus \{0\}] + 1$ we have $[R \setminus \{0\}] = -1$. Together with $0 = [(R \setminus \{0\})^2] = [R \setminus \{0\}]^2$ this yields $1 = 0$, so $K_0(K, \mathcal{L})$ is trivial. Write g_2 for the isomorphism

$$g_2 : (R \setminus \{0\})^2 \rightarrow (R \setminus \{0\})^2 \sqcup (R \setminus \{0\})^2$$

Now take the disjoint union of $R^{(1)} \times (R \setminus \{0\})$, $R \times (R \setminus \{0\})$ and $(R \setminus \{0\})^2$ inside R^2 in some way, meaning that we take disjoint isomorphic copies inside R^2 of the mentioned sets. Using the above isomorphisms g_1 and g_2 in a clever way on these disjoint copies, it is clear that we can remove one copy of $R \times (R \setminus \{0\})$ from R^2 and put one copy of $(R \setminus \{0\})^2$ back instead, hence we find an isomorphism from R^2 to itself minus a point. For details of this construction, we refer to the proof of [1], theorem 1. $\square$

4. THE PROOFS OF THEOREMS 1, 2, AND 3

Proof of theorem 1. Fix a field K as in the statement. Using Hensel's lemma, it is elementary to calculate for each n the numbers $r_n(K)$ and $s_n(K)$, and to find that, for n a prime number, $s_n(K)/r_n(K)$ is a power of n . Further, it is not difficult to check that taking n -th roots is definable. Therefore, the generator $\lambda_2 - 1$ of $H(K, \mathcal{L}_{\text{ring}})$ is uneven and $\lambda_3 - 1$ is even. This implies that $H(K, \mathcal{L}_{\text{ring}}) = \mathbb{Z}$.

We calculate explicitly for $K = \mathbb{Q}_p((t))$, for the other fields of the statement, the arguments are completely similar, although, notation can get more complicated.

Let v be the valuation on K into $\mathbb{Z} \times \mathbb{Z} \cup \{\infty\}$ with lexicographical order, determined by: $v(x) = (s, r)$ for $x = \sum_{i \geq s} a_i t^i$ with $a_i \in \mathbb{Q}_p$ and $a_s \neq 0$, the p -adic valuation of a_s being r . The valuation ring R is definable and can be described by

$$R = \{x \in K \mid 1 + tx^2 \in P_2(K) \text{ \& } 1 + px^2 \in P_2(K)\}$$

if $p \neq 2$ and by

$$R = \{x \in K \mid 1 + tx^3 \in P_3(K) \text{ \& } 1 + px^3 \in P_3(K)\}$$

if $p = 2$. Write M for the maximal ideal of R . Let $\text{ac} : K \rightarrow \mathbb{F}_p$ be the angular component $\text{ac}(x) = p^{-r} t^{-s} x \bmod M$ for nonzero x with $v(x) = (s, r)$. The set $R^{(1)} = \{x \in R \mid \text{ac}(x) = 1\}$ is definable since it is the union of the sets

$$p^i t^j P_{p-1}(K),$$

for $i, j = 0, \dots, p-2$.

Now we can use proposition 1, to find a $\mathcal{L}_{\text{ring}}$ -definable bijection $K \rightarrow K^\times$ and to find that $K_0(K, \mathcal{L}_{\text{ring}})$ is trivial. This proves the proposition for fields of the form $\mathbb{Q}_p((t))$. When L is a finite field extension of $\mathbb{Q}_p$ and K an iterated Laurent

series field over L , there are $\mathcal{L}_{\text{ring}}$ -formula's playing the role of φ in the obvious way and the reader can make the adaptations. $\square$

Proof of Theorem 2. Suppose for simplicity that K is the field $(\mathbb{F}_p((t_1)))((t_2))$, where p is a prime. The other cases are completely similar. Let v_2 be the valuation on K into $\mathbb{Z} \times \mathbb{Z}$ as in section 1.2; this is a valuation determined by: $v(x) = (s, r)$ for a Laurent series $\sum_{i \geq s} a_i t_2^i$ with $a_i \in \mathbb{F}_q((t_1))$ and $a_s \neq 0$ and $a_s = \sum_{i \geq r} b_i t_1^i$ with $b_r \neq 0$ and $b_i \in \mathbb{F}_q$. Write M for the maximal ideal with respect to v_2 . The valuation ring $R = \{x \mid v_2(x) \geq 0\}$ is $\mathcal{L}_{\text{ring}}$ -definable because of the following observation:

$$R = \{x \in K \mid 1 + t_2 x^2 \in P_2(K) \text{ \& } 1 + t_1 x^2 \in P_2(K)\},$$

if $\text{char}(K) \neq 2$ and

$$R = \{x \in K \mid 1 + t_2 x^3 \in P_3 \text{ \& } 1 + t_1 x^3 \in P_3\},$$

if $\text{char}(K) = 2$.

Let $\text{ac} : K \rightarrow \mathbb{F}_p$ be the angular component map $x \mapsto t_2^{-s} t_1^{-r} x \bmod M$ for nonzero x with $v_2(x) = (s, r)$. The set $R^{(1)} = \{x \in R \mid \text{ac}(x) = 1\}$ is definable since it is the union of the sets

$$t_1^j t_2^i P_{p-1}(K)$$

for $i, j = 0, \dots, p-2$. Now use proposition 2 to find that $K_0(K, \mathcal{L}_{\text{ring}})$ is trivial and to find a $\mathcal{L}_{\text{ring}}$ -definable bijection

$$g_3 : R^2 \rightarrow R^2 \setminus \{(0, 0)\}.$$

The following is a definable injection:

$$g_4 : R^2 \rightarrow R : (x, y) \mapsto x^p + t_1 y^p,$$

and thereby, we can define the $\mathcal{L}_{\text{ring}}$ -isomorphism

$$g_5 : R \rightarrow R \setminus \{0\} : x \mapsto \begin{cases} g_4 g_3(g_4^{-1}(x)) & \text{if } x \in g_4(R^2) \\ x & \text{else.} \end{cases}$$

This finishes the proof. $\square$

Proof of theorem 3. Let K be the field $\mathbb{R}((t_1)) \dots ((t_{n-1}))((t_n))$. Taking n -th roots is clearly $\mathcal{L}_{\text{ring}}$ -definable, and, using the notation of definition 1 we have that $\lambda_2(K, \mathcal{L}_{\text{ring}})$ is a power of two and $\lambda_3(K, \mathcal{L}_{\text{ring}})$ is a power of three. Therefore, $H(\mathbb{R}((t)), \mathcal{L}_{\text{ring}}) = \mathbb{Z}$. The existence of a $\mathcal{L}_{\text{Pas}}$ -definable bijection $K \rightarrow K^\times$ and the triviality of $K_0(K, \mathcal{L}_{\text{Pas}})$ are formal consequences of proposition 1, because $H(K, \mathcal{L}_{\text{ring}}) = \mathbb{Z}$, the value group clearly has a unique minimal strictly positive element and $R^{(1)}$ is $\mathcal{L}_{\text{Pas}}$ -definable. $\square$

Proof of theorem 4. Let K be $L((t_1)) \dots ((t_n))$ where L is a field of characteristic $p > 0$. The statement follows immediately from proposition 2 using the definable injection

$$R^2 \rightarrow R : (x, y) \mapsto x^p + t_1 y^p$$

as in the proof of theorem 2. $\square$

REFERENCES

1. Raf Cluckers and Deirdre Haskell, *Grothendieck rings of $\mathbb{Z}$ -valued fields*, Bulletin of Symbolic Logic **7** (2001), no. 2, 262–269.
2. J. Denef and F. Loeser, *Germes of arcs on singular algebraic varieties and motivic integration*, Inventiones Mathematicae **135** (1999), 201–232.
3. ———, *Definable sets, motives and p -adic integrals*, Journal of the American Mathematical Society **14** (2001), no. 2, 429–469.
4. J. Krajíček and T. Scanlon, *Combinatorics with definable sets: Euler characteristics and grothendieck rings*, Bulletin of Symbolic Logic **6** (2000), 311–330.
5. J. Pas, *Uniform p -adic cell decomposition and local zeta functions*, Journal für die reine und angewandte Mathematik **399** (1989), 137–172.
6. ———, *On the angular component map modulo p* , J. Symbolic Logic **55** (1990), 1125–1129.
7. Lou van den Dries, *Tame topology and o-minimal structures*, Lecture note series, vol. 248, Cambridge University Press, 1998.

DEPARTMENT OF MATHEMATICS, KATHOLIEKE UNIVERSITEIT LEUVEN, CELESTIJNENLAAN 200B,
B-3001 LEUVEN, BELGIUM

E-mail address: `raf.cluckers@wis.kuleuven.ac.be`

URL: `http://www.wis.kuleuven.ac.be/algebra/raf/`